



---

## **The Effect of William Flexion Exercise on Reducing Pain Intensity For Elderly with Low Back Pain**

**Amila<sup>1\*</sup>, Henny Syapitri<sup>2</sup>, Evariana Sembiring<sup>3</sup>**

<sup>1</sup> Departement of Medical Surgical Nursing, Sari Mutiara Indonesia University, Medan, Indonesia  
<sup>2,3</sup> Departement of Public Health, Sari Mutiara Indonesia University, Medan, Indonesia

---

### **Article info**

**Article history:**

Received; Agustus 15<sup>th</sup>, 2020

Revised: September 15<sup>th</sup>, 2020

Accepted: October 15<sup>th</sup>, 2020

---

**Correspondence author:**

Amila

E-mail: [mila\\_difa@yahoo.co.id](mailto:mila_difa@yahoo.co.id)

---

**DOI:**

<http://doi.org.10.35654/ijnhs.v4i1.374>

---

### **Abstract**

The elderly are a group of the population that has the potential to become vulnerable communities. In the elderly, the aging process can cause various physical problems. One of the issues that often arise in the elderly is low back pain (NPB). The purpose of this study is to evaluate the effect of the William Flexion Exercise on reducing the pain intensity for elderly with low back pain at Hisosu Binjai Resident Home, Indonesia. The research design is quasi-experimental with the pre and post-test control group approach in 2017. This study's sample consisted of 28 people in the intervention group and 28 people in the control group. The sampling technique utilized consecutive sampling. This study used The Pain Numerical Rating Scale to evaluate the level of pain intensity perceived by the patient. The data were analyzed by dependent and independent t-test with significance  $\alpha < 0.05$ . The dependent t-test found a significant difference in the low back pain before and after the intervention group intervention (p-value = 0.000). In contrast, there is no significant difference in the control group (p-value = 0.081). The result of the independent t-test found that there are significant differences in the low back pain between the intervention group and the control group. This study concludes that William's flexion exercise provided benefits to most of the participants in this study. Thus, it may be an effective technique to reduce pain intensity and increase the range of motions.

**Keywords:** aging exercise therapy; low back pain; flexion exercise

This is an Open Access article distributed under the terms of the Creative Commons Attribution 4.0 International License CC BY - 4.0



## INTRODUCTION

Low back pain (LBP), perhaps more accurately called lumbago or lumbosacral pain, occurs below the 12<sup>th</sup> rib and above the gluteal folds. LBP is one of the most common health problems all over the world (1,2). The lifetime prevalence of non-specific LBP may reach 80%, with the annual prevalence ranging between 25% and 60% in different ethnic groups (3). World Health Organization (WHO) reported that the world population aged over 60 years would increase from the current 841 million to two billion by 2050, turning chronic diseases and older people's welfare into new challenges for global public health (4-5).

Many risk factors caused low back pain, such as the worker who spent all their time by impressive duty, holder and lifted bulky goods, smoker, obesity, and activities. It was due to sit for a long time and influenced by the aging process (6). While most LBP causes among older adults are non-specific and self-limiting, seniors are prone to develop specific LBP pathologies and chronic LBP given their age-related physical and psychosocial changes (7). LBP often occurs in the elderly (8-9).

A variety of changes in the elderly occurred on a musculoskeletal system, including porous bones, enlargement of joints, shift a tendon, limited motion, depletion of intervertebral disc and muscular weakness aging process. The bone reached its peak at the age of 35 years. After it happened, bone losses thorough gradually. In the elderly, the structure of collagen less able to absorb energy. Therefore, muscle does, and power is also reduced. Loss occurs the number of muscle fiber due to atrofimiofibril with the replacement of tissue fibrous that start to develop in the life in the fourth (10). The lumbar spine structures involved in the development of low back pain are intervertebral disk cartilages, intervertebral joints, tendons, and muscles. When the sensory receptors in these structures receive nociceptive stimulation, they trigger a pain reaction in the pain sensation system, including both at the peripheral and the central levels. Ligament and muscles that surround the lumbar spine. Strain in these structures results in back pain.

Inappropriate posture, irregular movement of the lumbar vertebrae, and reduced or imbalanced muscle strength enhance the nociceptive stimulation. Motion restriction due to pain leads to intervertebral joints' contracture and the atrophy of the other lumbar spine structures, resulting in a vicious circle of pain (11).

The results indicate a high prevalence of LBP among elderly individuals, ranging from 21.7 to 75%. Furthermore, the majority of LBP is high in developed countries such as Canada (75%) (5), the United States (67%) (12), Sweden (49%) (13), China (39.2%) (14) and Japan (32%) (15). LBP occurs in 43% of both men and women, differing from the mean global prevalence, 31%. This finding was also confirmed in developing countries such as Brazil, where the majority was 33.6% to 68.3%. In other Brazilian studies, the small samples studied may have contributed to a high prevalence, and the samples may not have been representative of the study population (16).

There is still no epidemiology data about low back pain in Indonesia. Approximately 40% of Central Java society has over 65 years old whoever had low back pain. The Community Oriented Program for Control of Rheumatic Disease (COPCRD) of Indonesia showed that the prevalence of low back pain is about 18,2 % for men and 13,6 % for women. The incidence is based on patients visiting some hospitals in Indonesia, about between 3-17 % (17).

LBP recently was rated by the Global Burden of Disease Study as one of the seven health conditions that most affect the world's population. It is considered a debilitating health condition that affects people for the most significant number of years over a lifetime. LBP is a major cause of disability – affecting performance at work and general well-being. It is also associated with high treatment costs (18), (19).

Older adults are one of the communities that can become facilitate people so that need to create a physical and non-physical condition. More elderly who have low back pain handle it with resting, drink some traditional treatment or let it so. Older adults often get hurt on their low back if not taking muscle stretching as soon as possible by stretching so the joint will be small and resulting in pain. Reducing low back pain in patients is recommended with conservative therapy. By this exercise, therapy on low back cases had become a routine treatment for every practitioner in physiotherapy, but a nurse also can do the treatment. As role models in society, nurses have a significant role in obligating the pain through a non-pharmacology approach. Interventions that

include non-pharmacology approaches, such as giving good exercise (20).

Back exercise is one of the activities that give a great use. The back exercise routinely for a long time will actively progress the power of muscle, that called active stabilization. The progress of muscle power also affects the body's progress toward movement or loaded statistically and dynamically. There are many types of treatments for low back pain. These vary from conventional methods (physiotherapy, kinesiotherapy, massage) to modern techniques like ozone therapy. However, two protocols stand out due to their popularity, like the Williams and McKenzie protocols (21).

William Flexion Exercise is designed to reduce back pain by strengthening the muscles that reflect the lumbosacral spine, especially the abdominal and maximal gluteus muscles, and stretching the extensor muscle groups. The principle of exercise therapy using the William Flexion is to repair body posture, reduce lumbal hyperlordosis. It also decreases muscle spasms through relaxing effects, avoids stiff intervertebral joints, and checks the lousy posture (21-22). The back exercise was taken in a month (3 times in a week) showed a meaningful difference in the decrease of pain with the method of William Flexion (10).

Back exercise is an easy movement because it just has six moving and spends 15 – 20 minutes in every method. This method can be done three times a week (12). Besides easy to imitate, this exercise does not need sophisticated tools and special place, cheaper, and it can do by ourselves based on Standard Operating Procedure (SOP). Back exercise also gives influence toward the muscle bend progress, elastically bone and it can be reducing of low back pain.

Unlike other therapeutic methods, this method aims to make the patients as independent of the therapist as possible and thus capable of controlling their pain through postural care and the practice of specific exercises for their problem. It encourages patients to move the spine in a direction that is not harmful to their problem, thus avoiding movement restriction due to pain (12).

## **OBJECTIVE**

The study aimed to evaluate William Flexion's effectiveness in reducing pain intensity for elderly with low back pain in Hisosu Binjai Resident Home.

## **METHOD**

This research is a comparative analytic study using a quasi-experimental study with a pre and post-test control group approach. The samples of this study were recruited with consecutive sampling. Inclusion criteria include: never got same therapy, pain intensity before doing research was minimum to moderate, able to speak Indonesia, willing to followed research program entirely for four weeks, and low back pain did not accompany with neurological disturbance. Patients will be excluded if they have any contraindication to physical exercise include evidence of nerve root compromise (i.e., one or more moving, reflex, or sensation deficits). We also excluded the severe spinal pathology (e.g., fracture, tumor, inflammatory and infectious diseases), serious cardiovascular and metabolic diseases, previous back surgery. At the beginning and the end of the study, the participants were given the Numeric Pain Intensity Scale, which design to measure the pain of intensity scale.

This study used The Pain Numerical Rating Scale to evaluate the level of pain intensity perceived by the patient. The measurement for assessing pain using the Numeric Pain Intensity Scale (NPIS) has been translated into the Indonesian language. This NPIS was selected because this instrument was affordable and easy to administer in health services. The NPIS is a valid and reliable scale to measure pain intensity (24).



t-test), firstly, we examined the normality data by Shapiro Wilk Test. According to the result of the Shapiro Wilk test, data were normally distributed. The effectiveness of William Flexion in the elderly was analyzed using a Dependent and Independent sample t-test with significance  $\alpha < 0.05$ .

## RESULT

Based on table 1, the mean age of the elderly in the control group was 67.85 (SD = 4.160), and the mean age of the elderly in the intervention group was 67.29 (SD = 4.730).

Table 1. The Characteristics in Elderly with Low Back Pain (n = 28)

Variable	Group	Mean	SD	Min-Max	95% CI
	Intervention	67.29	4.730	61 -74	64.55– 70.02
	Control	67.85	4.160	60 -74	65.33 – 70.36

Based on the data in table 2, it can be seen that respondents in the control group had the same number of both males, i.e., 14 respondents (50%) and female, i.e., 14 respondents (50%). Meanwhile, most of the intervention group respondents were male, i.e. 16 respondents (57.14%), and the rest were female, i.e., 12 respondents (42.86%). Based on the lifestyle of smoking, most of the respondents in the control group is no smoker, i.e., 15 respondents (53.57%), and the rest were smokers was 13 respondents (46.43%). Meanwhile, in the intervention group is smoker, i.e., 15 respondents (53.57%), and the rest were no smoker, i.e., 13 respondents (46.43%).

Table 2. The Characteristics in Elderly with Low Back Pain (n=28)

Variables	Category	Intervention Group		Control Group	
		N	%	N	%
	Male	16	57.14	14	50.00
	Female	12	42.86	14	50.00
LifeStyle of Smoking					
	Smoker	15	53.57	13	46.43
	No Smoker	13	46.43	15	53.57

Table 3 shows that based on the statistical analysis using the dependent sample t-test in the intervention group (p-value is 0.0000) and the control group (p-value is 0.081) at  $\alpha = 0.05$ . It may indicate a significantly different result of the low back pain before and after receiving the intervention (p-value =0.000). In contrast, there is no significant difference in the control group (p-value = 0.081). The result of the independent t-test found that there are significant differences in the low back pain between the intervention group and the control group. It may indicate a significantly different result between low back pain in control and those in the intervention group.

Table 3. The William Flexion Exercise in Elderly with Low Back Pain (n = 28)

Group	Pain Intensity	Mean	SD	SE	P-value
Intervention	Pretest	5.08	0.82	0.239	0.000
	Post-test	3.23	0.83	0.231	
Control	Pre test	4.79	0.76	0.128	0.081
	Post-test	4.26	0.64	0.116	

$\alpha = 0.05$

## DISCUSSION

### The Characteristics in Elderly with Low Back Pain

The findings of this study were consistent with previous studies. Among changes in lumbar spine structures, age-related degeneration of intervertebral disk cartilages and intervertebral joints are common causes of low back pain. Aging is a well-known risk factor of LBP as degenerative changes in the spine and disc. At the age of 60-year-old, muscle power dropping by 20%, and risk complaints muscle will increase. Increasing age will bellow the function of the human body system, including the musculoskeletal system. This will lead to a rise in musculoskeletal complaints, including low back pain (11). A Spanish study found that the prevalence rates of chronic LPB among females and males aged 65 years or older were 24,2% and 12,3% (27), respectively, while an Israeli study documented that the prevalence of chronic LBP in people aged years was as high as 58% (28).

The age-related changes in neuroplasticity may decrease pain tolerance in older adults. Compared to younger adults, older people tend to show more rapid temporal summation of noxious heat stimuli in their central nervous system. Similarly, older adults display a prolonged period of capsaicin-induced hyperalgesia. It may lead to persistent pain sensitization and sluggish resolution of neoplastic change. Notably, central pain processing can be further complicated by dementia-related neurodegeneration.

This study, inconsistent with Cho et al., explained that the lifetime prevalence of LBP was 61.3%, with women having a higher prevalence (29). The point and 6-month prevalence were also higher among women. Some studies have shown that males are at greater risk for low back pain, while other studies suggest that females are more likely to develop this type of pain. Women who have had two or more pregnancies have a higher risk of developing low back pain. Jimenez-Sanchez and coworkers (2017) estimated that women were two times more likely to develop chronic LBP than men. The higher prevalence of chronic pain in females may be attributed to complex bio-psychosocial. Further, women commonly have a higher number of concomitant chronic diseases (e.g., osteoporosis, osteopenia, and osteoarthritis), which are known to be risk factors for developing chronic LBP and psychological distress in older adults (27), (30)

This study, consistent with Miranda et al. (2008), showed an association between smoking and LBP among over 50 years old. Smoking is related to a higher propensity for LBP in older people (27). Also, another study showed an association between smoking and decreased bone mineral density in women. Former smokers had a higher prevalence of low back pain than never smokers, but a lower prevalence of low back pain than current smokers. In cohort studies, both former (OR 1.32, 95% CI, 0.99-1.77) and current (OR 1.31, 95% CI, 1.11-1.55) smokers had an increased incidence of low back pain compared with never smokers. If you are a smoker, you are 2.7 times more likely to develop lower back pain than if you didn't smoke (31).

The researchers found that women who smoked complained more often of back pain than women who didn't smoke in both groups, not just the scoliosis group (32). It's not entirely understood how cigarette smoking affects the back. One theory is that nicotine causes vasoconstriction, or narrowing of the blood vessels, that provide nutrition to the discs' cells. If the nutrients can't reach the cells, this leads to malnutrition of the disc, and they can become damaged more easily. Malnourished tissues also can't heal themselves as quickly or as well as healthier, nourished cells. Nicotine is also known to thicken the walls of the blood vessels. This has the same effect of narrowing the blood vessels, slowing down blood flow. One of the by-products of cigarettes is carbon monoxide, an extremely poisonous gas. Carbon monoxide is also blamed for the increase in lower back pain. When you smoke, the carbon monoxide attaches itself to your hemoglobin, the part of your blood cells that carry oxygen to the tissues throughout your body. This burden on the

hemoglobin takes up space, keeping much-needed oxygen from reaching the discs in your back. Like vasoconstriction, this causes malnutrition to the cells (31-32).

### **The Effect of William Flexion Exercise on Reducing the Pain Intensity for Elderly with Low Back Pain**

Results of observation based on research variables found  $p = 0.000$  ( $p < 0.05$ ), which means there is a difference in the low back pain between the intervention and control groups. The study was consistent with (21) that there were significant effects of William's flexion exercise in inadequate back pain management. Pain intensity was significantly reduced. This finding supports the result of a study conducted by Sukmajaya, Alkaff, Oen, Sukmajaya (2020), showing that there was a significant ODI/ Oswestry Disability Index/ difference between pre-and-post-WFE implementation ( $31.05 \hat{\pm} 17.40$  vs.  $14.10 \hat{\pm} 11.78$ ,  $p = 0.019$ ) (33). There was no significant difference in ODI reduction between geriatric and non-geriatric participants ( $p = 0.24$ ). It was consistent with Yundari & Mas (2018) explained that William's Flexion Exercise given to the woodcarvers significantly reduce the pain intensity ( $p$ -value = 0.000) (34).

About William Flexion, results are compatible with previous studies, which all of them known this protocol as an effective method. The William Flexion method involves repeated movements or sustained positions and has an educational component to minimize pain and disability and improve spinal mobility (23). The primary benefit is the opening of the intervertebral foramen, the stretching of ligamentous structures, and the distraction of the apophyseal joints. Previous studies demonstrated that active therapy could improve functional ability and reduce pain when it has both exercise and patient collaboration (35) explicitly.

The average decrease of low back pain caused flexion back muscle strengthens through William Flexion exercise to reduce the pain. It has related to muscle work. The more muscle strength, the more muscle is having a contraction, and on the contrary. The muscles do their function with the couple because when agonist muscle contraction, antagonist muscle in another way in relaxation. If this has not happened, the two muscles will drive each other blocking the movement and resulting in pain (21 22).

Flexion exercise can help reduce pain by reducing intradiscal pressure (21, 35-36) Someone who does the training,  $\beta$ -endorphin will be captured by receptors in the hypothalamus and the limbic system regulating emotions. Increased  $\beta$ -endorphin proved to be closely related to decreased pain, increased memory, improved libido, sexual ability, blood pressure, and respiration (37). This flexion exercise aims to reduce the body's pressure on the articular weight-bearing stress and stretch the muscles and fascia (increase the soft tissue extensibility) in the dorsolumbar region, useful for correcting the wrong posture. This flexion exercise also improves stability in the lumbar region by actively training the abdominal muscles, gluteus Maximus and hamstring (34).

## **CONCLUSION**

William Flexion exercise in elderly with low-back pain was effective in reducing pain. The results of this study will contribute to better management of this population. Also, compared to the other back exercises, include Mc. Kenzie exercise of elderly is recommended in Hisosu Binjai Resident Home.

## **ACKNOWLEDGEMENTS**

The authors are grateful to all the elderly at the Hisosu Binjai Resident Home, North Sumatera, for their cooperation during data collection.

## **REFERENCES**

- (1) L S, Hanifa S. Prevalence and risk factors of low back pain among nurses in a typical Nigerian hospital. *Afr Health Sci.* 2010;10(101):26–30.
- (2) Keriri HM, Keriri H. Prevalence and risk factors of low back pain among nurses in operating rooms, Taif, Saudi Arabia. *Am J Res Commun* www.usa-journals.com Keriri. 2013;1(11):45–70.
- (3) Mcbeth J, Jones K. *Epidemiology of chronic musculoskeletal pain.* Elsevier. 2007;21(3):403–25.
- (4) Cassidy JD, Co P, Carroll LJ, Kristman V. Incidence and Course of Low Back Pain Episodes in the

- General Population. 2005;30(24):2817–23.
- (5) Hoy D, March L, Brooks P, Blyth F, Woolf A, Bain C, et al. The global burden of low back pain : estimates from the Global Burden of Disease 2010 study. *BMJ J.* 2014;73(6):968–74.
  - (6) Hoogendoorn WE, van Poppel MN, Bongers PM, Koes BW, Bouter LM. Systematic review of psychosocial factors at work and private life as risk factors for back pain. *Spine (Phila Pa 1976).* 2000;25(16):2114–25.
  - (7) Wong AYL, Karppinen J, Samartzis D. Low back pain in older adults : risk factors, management options, and future directions. 2017;1–23.
  - (8) Schild von Spannenberg S, Jones GT, Macfarlane GJ. The evidence base for managing older persons with low back pain. *Br J Pain.* 2012;6(4):166–9.
  - (9) Gheno R, Cepparo JM, Rosca CE, Cotten A. Musculoskeletal Disorders in the Elderly. 2012;2(3).
  - (10) Freemont AJ, Hoyland JA. Morphology, mechanisms, and pathology of musculoskeletal aging. *J Pathol.* 2007;211(2):252–9.
  - (11) Yamamoto H. Low Back Pain Due to Degenerative Disease in Elderly Patients. *Japan Med Assoc J.* 2003;128(12):433–8.
  - (12) Marshall M, Litwack-Harrison S, Makris E, Kado DM, Cawthon PM, Deyo RA, et al. A Prospective Study of Back Pain and Risk of Falls Among Older Community-dwelling Men. *Journals Gerontol.* 2017;72(9):1264–9.
  - (13) Kherad M. Risk factors for low back pain and sciatica in elderly men — the MrOS Sweden study. 2014;64–71.
  - (14) Hjermstad MJ, Fayers PM, Haugen DF, Caraceni A, Hanks GW, Med D, et al. Studies Comparing Numerical Rating Scales, Verbal Rating Scales, and Visual Analogue Scales for Assessment of Pain Intensity in Adults : A Systematic Literature Review. *J Pain Symptom Manage [Internet].* 2011;41(6):1073–93. Available from: <http://dx.doi.org/10.1016/j.jpainsymman.2010.08.016>
  - (15) Obuke YK, Yasuyo ABE, Zhaojia YE, Onda SH, Ta MTOMI, Saki MO, et al. Association of Age, Obesity, Joint Pain, and Chewing Ability with Chair Stand Difficulty among Community-dwelling Elderly People in Nagasaki, Japan. 2009;(0):4–7.
  - (16) Palma R, De Conti MHS, Quintino NM, Nuevo GMA. FUNCTIONAL CAPACITY AND ITS ASSOCIATED FACTORS IN THE ELDERLY WITH LOW BACK PAIN. 2014;22(6):295–9.
  - (17) Katz JN. Lumbar Disc Disorders and Low-Back Pain: Socioeconomic Factors and Consequences. *J Bone Jt Surg.* 2006;88(suppl\_2):21.
  - (18) Murray CJL, Lopez AD. Measuring the Global Burden of Disease. *N Engl J Med.* 2013;369(5):448–57.
  - (19) Nedergaard A, Henriksen K, Karsdal MA, Christiansen C. Best Practice & Research Clinical Obstetrics and Gynaecology Musculoskeletal aging and primary prevention. *Best Pract Res Clin Obstet Gynaecol [Internet].* 2013;1–16. Available from: <http://dx.doi.org/10.1016/j.bpobgyn.2013.06.001>
  - (20) Hayden JA, Cartwright J, van Tulder MW MA. Exercise therapy for chronic low back pain (protocol). *Cochrane Database Syst Rev.* 2012;(4): Art. No.: CD009790.
  - (21) Kumar M, Educational MGR. Effectiveness of William ' S Flexion Exercise in the Management of Low. 2016;(February).
  - (22) Gupta S. A COMPARISON BETWEEN McKENZIE EXTENSIONS EXERCISES VERSUS WILLIAM'S FLEXION EXERCISES FOR LOW BACK PAIN IN B.PT. STUDENTS. Vol. 3, *Indian Journal of Physical Therapy.* 2015.
  - (23) Ponte DJ, J. JG, Kent BE. A preliminary report on the use of the McKenzie protocol vs. Williams protocol in the treatment of LBP. *J Orthop Sports Phys Ther.* 1984;6(10):130–7.
  - (24) Hawker GA, Mian S, Kendzerska T, French M. Measures of adult pain: Visual Analog Scale for Pain (VAS Pain), Numeric Rating Scale for Pain (NRS Pain), McGill Pain Questionnaire (MPQ), Short-Form McGill Pain Questionnaire (SF-MPQ), Chronic Pain Grade Scale (CPGS), Short Form-36 Bodily Pain Scale (SF. *Arthritis Care Res.* 2011;63(SUPPL. 11):1–26.
  - (25) Childs JD, Piva SR, Fritz JM. Responsiveness of the numeric pain rating scale in patients with low back pain. *Spine (Phila Pa 1976).* 2005;30(11):1331–4.
  - (26) Voinea A, Iacobini A. WILLIAMS ' PROGRAM FOR LOW BACK PAIN. VI(4):210–4.

- (27) Carrasco-garrido P, Jiménez-sánchez S, Fernández-de-las-pe C, Hernández-barrera V, Alonso-blanco C, Palacios-ce D, et al. Prevalence of chronic head , neck and low back pain and associated factors in women residing in the Autonomous Region of Madrid ( Spain ). 2012;26(6):534–40.
- (28) Jacobs JM, Hammerman-Rozenberg R, Cohen A, Stessman J. Chronic Back Pain Among the Elderly : Prevalence, Associations, and Predictors. 2006;31(7):203–7.
- (29) Cho NH, Jung YO, Lim SH, Chung C-K, Kim HA. The Prevalence and Risk Factors of Low Back Pain in Rural Community Residents of Korea. *Spine (Phila Pa 1976)*. 2012;37(24):2001–10.
- (30) Kim W, Jin YS, Lee CS, Hwang CJ, Lee SY, Chung SG, et al. Relationship Between the Type and Amount of Physical Activity and Low Back Pain in Koreans Aged 50 Years Older. *PM&R [Internet]*. 2014;(May). Available from: <http://dx.doi.org/10.1016/j.pmrj.2014.04.009>
- (31) Alkherayf F, Agbi C. Cigarette smoking and chronic low back pain in the adult population. Vol. 32, *Clinical and investigative medicine. Médecine clinique et experimentale*. 2009. p. E360-7.
- (32) Shiri R, Falah-Hassani K. The Effect of Smoking on the Risk of Sciatica: A Meta-analysis. *Am J Med*. 2016;129(1):64-73.e20.
- (33) Sukmajaya WP, Alkaff FF, Oen A, Sukmajaya AC. Williams Flexion Exercise for Low Back Pain : A Possible Implementation in Rural Areas. *Maced J Med Sci*. 2020;8(B):1–5.
- (34) Dalem AAI, Yundari H, Puspita P, Mas W. Proceedings of International Conference on Applied Science and Health ICASH-A44 EFFECTIVENESS OF WILLIAM FLEXION EXERCISE TO REDUCE PAIN INTENSITY ON LOW BACK PAIN ( LBP ) OF WOODCARVERS IN BALI, INDONESIA Proceedings of International Conference on Appli. 2018;456(3):305–11.
- (35) Ghiasi F, Mehraeen M. The effect of William’s exercise on non-specific and chronic referral low back pain. *Q Kermanshah Univ Med Sci*. 2008;12(4):330–42.
- (36) MOLDOVAN M. Therapeutic Considerations and Recovery in Low Back Pain: Williams vs. McKenzie. *Timisoara Phys Educ Rehabil J*. 2012;5(9):58–64.
- (37) Siswantoyo. Perubahan Kadar Beta Endorphin Akibat Latihan Olahraga Pernafasan (sebuah Kajian Psikoneuroendokrinologi pada Aktivitas Fisik). 2010. p. 157–62.